

December 6, 2005

Dear Healthcare Professional,

Thank you for your interest in the safety profile of GAMMAGARD LIQUID [Immune Globulin Intravenous (Human)] 10% with respect to an outbreak of the *SARS virus*. GAMMAGARD LIQUID is indicated for the treatment of primary immunodeficiency disorders associated with defects in humoral immunity. These include but are not limited to congenital X-linked agammaglobulinemia, common variable immunodeficiency, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies.<sup>1</sup>

In response to the worldwide outbreak of Severe Acute Respiratory Syndrome (SARS), Baxter would like to assure you that our medical and scientific staff has monitored the situation closely and has followed guidance from the US Centers for Disease Control and Prevention (CDC), the US Food and Drug Administration (FDA), and other safety agencies, as they have been issued<sup>2,3</sup>. As part of our constant commitment to safety, we have vigilantly monitored new diseases and revise our processes to ensure the utmost in therapy safety. Our current comprehensive multi-step safety programs, from donor screening to plasma processing, helps protect against the transmission of the SARS virus by our plasma-derived therapeutics, including GAMMAGARD LIQUID.

According to the US Food and Drug Administration (FDA) and World Health Organization (WHO), to this date there have been no reports of SARS transmission via plasma-derived therapeutics, yet persons with SARS could potentially have virus in their blood early in infection without displaying any symptoms. For this reason, our current standard operating procedures (SOPs) **exclude ill individuals** from donating plasma. On May 19, 2003 Baxter BioScience and BioLife Plasma Services, our affiliated plasma collection system, implemented an SOP change to include the FDA recommendations for the industry in cases of SARS or suspected SARS. Among these procedures were: Donor Interview Questions, Donor Post-donation Information and Product Look Back Investigation. On March 15, 2004, these SOP changes were discontinued in accordance with the FDA guidelines that recommended that SARS donor questions be discontinued ninety days after CDC had lifted all travel alerts for SARS affected areas.

The U.S. FDA has stated in Guidance for Industry that SARS is unlikely to be transmitted through therapeutics that are processed from plasma. The FDA believes that licensed plasma derivatives undergo additional viral reduction procedures that are validated to be effective against lipid-enveloped RNA viruses such as coronavirus.

In addition to donor screening, there are various virus inactivation/removal steps utilized in the preparation of GAMMAGARD LIQUID that are validated to be effective against lipid-enveloped RNA viruses such as coronaviruses. A novel member of the *Coronaviridae* family of viruses has been identified as the causative agent of SARS.

Baxter has performed extensive virus inactivation/removal studies on a variety of lipid-enveloped viruses with procedures such as organic solvent/detergent (S/D) treatment, the dedicated viral inactivation step employed in GAMMAGARD LIQUID processing. The results from our S/D inactivation studies have demonstrated that this class of virus is rapidly and effectively inactivated to a high level, and this step is extremely robust for clearance of lipid-enveloped viruses. Generally, the efficacy of the S/D treatment was also confirmed by a recent publication<sup>4</sup> that showed complete inactivation of SARS within one minute.

In addition to inactivation by the S/D treatment, enveloped viruses can also be removed during various steps of purification such as the Cohn-Oncley fractionation, 35 nm nanofiltration and low pH treatment at elevated temperature.

In conclusion, Baxter has measures in place to help prevent plasma donations containing SARS from entering the plasma purification process, as well as in-process controls and procedures which greatly minimize the risk of any possible transmission of the virus. As always, Baxter will continue to monitor the situation closely and take necessary actions if any additional safety concerns become apparent.

For more information on our plasma collection process or the viral inactivation/removal steps employed during the processing of GAMMAGARD LIQUID, please feel free to contact Baxter BioScience, Medical Affairs department, at **1-866-424-6724**.

Sincerely,

Richard I. Schiff MD, Ph.D.  
Medical Director of Medical Affairs  
Baxter BioScience

#### **References:**

1. GAMMAGARD LIQUID Package Insert, 2005  
[http://www.baxter.com/products/biopharmaceuticals/antibody\\_therapies/index.html](http://www.baxter.com/products/biopharmaceuticals/antibody_therapies/index.html)
2. <http://www.fda.gov/bbs/topics/ANSWERS/2003/ANS01215.html>
3. [http://www.fda.gov/fdac/features/2003/403\\_sars.html](http://www.fda.gov/fdac/features/2003/403_sars.html)
4. Rabenau HF, Biesert L, Schmidt T, Bauer G, Cinatl J, Doerr HW, SARS-coronavirus (SARS-CoV) and the safety of a solvent/detergent (S/D) treated immunoglobulin preparation, *Biologicals*. 2005; **33(2)**:95-9

BAXTER and GAMMAGARD are trademarks of Baxter International Inc. and are registered in the U.S. Patent and Trademark Office.

#### **Important Safety Information**

GAMMAGARD LIQUID is contraindicated in patients with known anaphylactic or severe hypersensitivity responses to Immune Globulin (Human). Patients with severe selective IgA deficiency (IgA < 0.05 g/L) may develop anti-IgA antibodies that can result in a severe anaphylactic reaction.

**Immune Globulin Intravenous (Human) products have been reported to be associated with renal dysfunction, acute renal failure, osmotic nephrosis, and death. Patients predisposed to acute renal failure include patients with any degree of pre-existing renal insufficiency, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs. Especially in such patients, IGIV products should be administered at the minimum concentration available and the minimum rate of infusion practicable. While these reports of renal dysfunction and acute renal failure have been associated with the use of many of the licensed IGIV products, those containing sucrose as a stabilizer accounted for a disproportionate share of the total number. Glycine, an amino acid, is used as a stabilizer. GAMMAGARD LIQUID does not contain sucrose.**

As with all plasma-derived therapeutics, the potential to transmit infectious agents (e.g. viruses) cannot be totally eliminated.

Thrombotic events have been reported in association with IGIV. Patients at risk may include those with a history of atherosclerosis, multiple cardiovascular risk factors, advanced age, impaired cardiac output, and/or known or suspected hyperviscosity, hypercoagulable disorders, and prolonged periods of immobilization.

Aseptic meningitis syndrome (AMS) has been reported to occur infrequently in association with IGIV treatment. Discontinuation of IGIV treatment has resulted in remission of AMS within several days without sequelae.

IGIV products can contain blood group antibodies that may cause a positive direct antiglobulin reaction and, rarely, hemolysis.

Components used in the packaging of this product are latex-free.

Various mild and moderate reactions, such as headache, fever, fatigue, chills, flushing, dizziness, urticaria, wheezing or chest tightness, nausea, vomiting, rigors, back pain, chest pain, muscle cramps, and changes in blood pressure may occur with infusions of Immune Globulin Intravenous (Human).

Please visit [www.gammagardliquid.com/pdf/gamliquid\\_PI.pdf](http://www.gammagardliquid.com/pdf/gamliquid_PI.pdf) for GAMMAGARD LIQUID Prescribing Information for full prescribing details

---

shortened half-life of IgG,<sup>14,15,16,17</sup>

In contrast, however, pharmacokinetic studies in immunodeficient patients are based on the decline of IgG concentrations following infusions of large quantities of gammaglobulin. In such trials, investigators have reported uniformly prolonged half-lives of 26 – 35 days.<sup>16,18,19,20,21,22</sup>

Pharmacokinetic parameters for GAMMAGARD LIQUID were determined from total IgG levels following the fourth infusion. A total of 61 subjects were enrolled and treated. Of these, 57 had sufficient pharmacokinetic data to be included in the dataset. Pharmacokinetic parameters are presented in Table 4.

**Table 4: Summary of Pharmacokinetic Parameters in 57 Subjects**

Parameter	Median	95% Confidence Interval
Elimination Half-Life (T <sub>1/2</sub> days)	35	(31, 42)
AUC <sub>0-24h</sub> (mg-days/dL)	29139	(27494, 30490)
C <sub>max</sub> (Peak, mg/dL)	2050	(1980, 2200)
C <sub>min</sub> (Trough, mg/dL)	1030	(939, 1110)
Incremental recovery (mg/dL)/(mg/kg)	2.3	(2.2, 2.6)

Abbreviations: AUC= area under the curve;

C<sub>max</sub>=maximum concentration; C<sub>min</sub>= minimum concentration

Median IgG trough levels were maintained between 960-1120 mg/dL. These dosing regimens maintained serum trough IgG levels considerably above 450 mg/dL, which is consistent with levels considered to be effective in the treatment of patients with Primary Immunodeficiency.<sup>23,24</sup> The elimination half-life of GAMMAGARD LIQUID of 35 days was similar to the half-lives reported for other IGIV products.<sup>13,14,15,17,25,26</sup>

## INDICATIONS AND USAGE

### Primary Immunodeficiency

GAMMAGARD LIQUID is indicated for the treatment of primary immunodeficiency disorders associated with defects in humoral immunity. These include but are not limited to congenital X-linked agammaglobulinemia, common variable immunodeficiency, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies.<sup>15,22</sup>

## CONTRAINDICATIONS

GAMMAGARD LIQUID is contraindicated in patients with known anaphylactic or severe hypersensitivity responses to Immune Globulin (Human).

Patients with severe selective IgA deficiency (IgA < 0.05 g/L) may develop anti-IgA antibodies that can result in a severe anaphylactic reaction. Anaphylaxis can occur using GAMMAGARD LIQUID even though it contains low amounts of IgA (average concentration of 37µg/mL). These patients should be treated only if their IgA deficiency is associated with an immune deficiency for which therapy with intravenous immune globulin is clearly indicated. Such patients should only receive intravenous immune globulin with utmost caution and in a setting where supportive care is available for treating life-threatening reactions.

## WARNINGS

**Immune Globulin Intravenous (Human) products have been reported to be associated with renal dysfunction, acute renal failure, osmotic nephrosis, and death.<sup>27</sup> Patients predisposed to acute renal failure include patients with any degree of pre-existing renal insufficiency, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs. Especially in such patients, IGIV products should be administered at the minimum concentration available and the minimum rate of infusion practicable. While these reports of renal dysfunction and acute renal failure have been associated with the use of many of the licensed IGIV products, those containing sucrose as a stabilizer accounted for a disproportionate share of the total number.**

**Glycine, an amino acid, is used as a stabilizer. GAMMAGARD LIQUID does not contain sucrose.**

**See PRECAUTIONS and DOSAGE AND ADMINISTRATION sections for important information intended to reduce the risk of acute renal failure.**

Immune Globulin Intravenous (Human), 10% is made from human plasma. Products made from human plasma may contain infectious agents, such as viruses, that can cause disease. The risk that such products will transmit an infectious agent has been reduced by screening plasma donors for prior exposure to certain viruses, by testing for the presence of certain current virus infections, and by inactivating and/or removing certain viruses (see DESCRIPTION). Despite these measures, such products can still potentially transmit disease. Because this product is made from human blood, it may carry a risk of transmitting infectious agents, e.g., viruses and theoretically, the Creutzfeldt-Jakob disease (CJD) agent. ALL infections thought by a physician possibly to have been transmitted by this product should be reported by the physician or other healthcare provider to Baxter Healthcare Corporation, at 1-800-423-2862 (in the U.S.). The physician should discuss the risks and benefits of this product with the patient.

GAMMAGARD LIQUID should only be administered intravenously. Other routes of administration have not been evaluated.

Immediate anaphylactic and hypersensitivity reactions are a remote possibility. Epinephrine and antihistamines should be available for treatment of any acute anaphylactoid reactions.

## PRECAUTIONS

### General

Some viruses, such as B19V (formerly known as Parvovirus B19) or Hepatitis A, are particularly difficult to remove or inactivate. B19V most seriously affects pregnant women, or immune-compromised individuals. Symptoms of B19V infection include fever, drowsiness, chills and runny nose followed about two weeks later by a rash and joint pain. Evidence of Hepatitis A may include several days to weeks of poor appetite, tiredness, and low-grade fever followed by nausea, vomiting and abdominal pain. Dark urine and a yellowed complexion are also common symptoms. Patients should be encouraged to consult their physician if such symptoms appear.

Components used in the packaging of this product are latex-free.

### Renal Function

Periodic monitoring of renal function tests and urine output is particularly important in patients judged to have a potential increased risk for developing acute renal failure. Assure that patients are not volume depleted prior to the initiation of infusion of GAMMAGARD LIQUID. Renal function, including measurement of blood urea nitrogen (BUN)/serum creatinine, should be assessed prior to the initial infusion of IGIV products and again at appropriate intervals thereafter. If renal function deteriorates, discontinuation of the product should be considered.

For patients judged to be at risk of developing renal dysfunction, it may be prudent to reduce the rate of infusion to less than 3.3 mg IgG/kg/min (<2 mL/kg/hr).

### Hemolysis

IGIV products can contain blood group antibodies which may act as hemolysins and induce *in vivo* coating of red blood cells with immunoglobulin, causing a positive direct antiglobulin reaction and, rarely, hemolysis.<sup>28,29,30</sup> Hemolytic anemia can develop subsequent to IGIV therapy due to enhanced red blood cells (RBC) sequestration (see ADVERSE REACTIONS).<sup>31</sup> IGIV recipients should be monitored for clinical signs and symptoms of hemolysis (see PRECAUTIONS: Laboratory Tests).

## Transfusion-Related Acute Lung Injury (TRALI)

There have been reports of noncardiogenic pulmonary edema (Transfusion Related Acute Lung Injury [TRALI]) in patients administered IGIV.<sup>32</sup> TRALI is characterized by severe respiratory distress, pulmonary edema, hypoxemia, normal left ventricular function, and fever, and typically occurs within 1-6 hours after transfusion. Patients with TRALI may be managed using oxygen therapy with adequate ventilatory support.

IGIV recipients should be monitored for pulmonary adverse reactions. If TRALI is suspected, appropriate tests should be performed for the presence of anti-neutrophil antibodies in both the product and patient serum (see PRECAUTIONS: Laboratory Tests).

## Thrombotic Events

Thrombotic events have been reported in association with IGIV (see ADVERSE REACTIONS).<sup>33,34,35,36,37,38,39,40,41</sup> Patients at risk may include those with a history of atherosclerosis, multiple cardiovascular risk factors, advanced age, impaired cardiac output, and/or known or suspected hyperviscosity, hypercoagulable disorders and prolonged periods of immobilization. The potential risks and benefits of IGIV should be weighed against those of alternative therapies for all patients for whom IGIV administration is being considered. Baseline assessment of blood viscosity should be considered in patients at risk for hyperviscosity, including those with cryoglobulins, fasting chylomicronemia/markedly high triacylglycerols (triglycerides), or monoclonal gammopathies (see PRECAUTIONS: Laboratory Tests).

## Aseptic Meningitis Syndrome

An aseptic meningitis syndrome (AMS) has been reported to occur infrequently in association with IGIV treatment. Discontinuation of IGIV treatment has resulted in remission of AMS within several days without sequelae. The syndrome usually begins within several hours to two days following IGIV treatment. It is characterized by symptoms and signs including severe headache, nuchal rigidity, drowsiness, fever, photophobia, painful eye movements, and nausea and vomiting. Cerebrospinal fluid (CSF) studies are frequently positive with pleocytosis up to several thousand cells per cubic mm, predominantly from the granulocytic series, and elevated protein levels up to several hundred mg/dL. Patients exhibiting such symptoms and signs should receive a thorough neurological examination, including CSF studies, to rule out other causes of meningitis. AMS may occur more frequently in association with high dose (2 g/kg) IGIV treatment.

## Laboratory Tests

If signs and/or symptoms of hemolysis are present after IGIV infusion, appropriate confirmatory laboratory testing should be done [see PRECAUTIONS].

If TRALI is suspected, appropriate tests should be performed for the presence of anti-neutrophil antibodies in both the product and patient serum [see PRECAUTIONS].

Because of the potentially increased risk of thrombosis, baseline assessment of blood viscosity should be considered in patients at risk for hyperviscosity, including those with cryoglobulins, fasting chylomicronemia/markedly high triacylglycerols (triglycerides), or monoclonal gammopathies [see PRECAUTIONS].

## Information For Patients

Patients should be instructed to immediately report symptoms of decreased urine output, sudden weight gain, fluid retention/edema, and/or shortness of breath (which may suggest kidney damage) to their physicians.

## Drug Interactions

See DOSAGE AND ADMINISTRATION section.

## Pregnancy Category C

Animal reproduction studies have not been conducted with GAMMAGARD LIQUID. It is also not known whether GAMMAGARD LIQUID can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. GAMMAGARD LIQUID should be given to a pregnant woman only if clearly indicated. Maternally administered IGIV products have been shown to cross the placenta, increasingly after 30 weeks gestation.<sup>42,43,44</sup>

## Use in Pediatrics

The safety and efficacy of GAMMAGARD LIQUID has not been evaluated in neonates or infants.

## ADVERSE REACTIONS

### General

Various mild and moderate reactions, such as headache, fever, fatigue, chills, flushing, dizziness, urticaria, wheezing or chest tightness, nausea, vomiting, rigors, back pain, chest pain, muscle cramps, and changes in blood pressure may occur with infusions of Immune Globulin Intravenous (Human). In general, reported adverse reactions to GAMMAGARD LIQUID in patients with Primary Immunodeficiency are similar in kind and frequency to those observed with other IGIV products. Slowing or stopping the infusion usually allows the symptoms to disappear promptly. Although hypersensitivity reactions have not been reported in the clinical studies with GAMMAGARD LIQUID immediate anaphylactic and hypersensitivity reactions are a remote possibility. Epinephrine and antihistamines should be available for treatment of any acute anaphylactic reactions (see WARNINGS).

### Clinical Study

Adverse experiences were examined among a total of 61 enrolled subjects with Primary Immunodeficiency who received at least one infusion of GAMMAGARD LIQUID during the Phase 3 multicenter clinical study. For this study, temporally associated adverse events are defined by the FDA as those occurring during or within 72 hours of completion of an infusion. Adverse drug reactions (ADR's) are those adverse events that were deemed by the investigators as causally related to the infusion of GAMMAGARD LIQUID.

Of all adverse experiences, 15 events in 8 subjects were serious. Two serious events, two episodes of aseptic meningitis in one patient, were deemed to be possibly related to the infusion of GAMMAGARD LIQUID.

Among the 896 non-serious adverse experiences, 258 were judged by the investigator to be possibly or probably related to the infusion of GAMMAGARD LIQUID. Of these, 136 were mild, 106 were moderate, and 16 were severe. All of the severe non-serious adverse experiences were transient, did not lead to hospitalization, and resolved without complication. One subject withdrew from the study due to a non-serious adverse experience (papular rash).

Of the 345 temporally related adverse experiences, those occurring in > 5% of subjects are shown in Table 5. Of these events, only headache occurred in association with more than 5% of infusions. All events were expected based on past experiences with intravenous gammaglobulin products.

Event	By Infusion		By Subject	
	Number	Percentage	Number	Percentage
Headache	57	6.90	22	36.1
Fever	19	2.30	13	21.3
Fatigue	18	2.18	10	16.4
Vomiting	10	1.21	9	14.8
Chills	14	1.69	8	13.1
Infusion site events	8	0.97	8	13.1
Nausea	9	1.09	6	9.8
Dizziness	7	0.85	6	9.8
Pain in Extremity	7	0.85	5	8.2
Diarrhea	7	0.85	5	8.2
Cough	5	0.61	5	8.2
Pruritus	5	0.61	4	6.5
Pharyngeal Pain	5	0.61	4	6.5

\* Excluding Infections

The majority (227/258) of the non-serious adverse experiences deemed related to study product were considered expected based on previous experience with IGIV products and 31 were considered unexpected. In virtually every case, these unexpected events were either consistent with the subject's specific type of immunodeficiency or with the subject's medical history prior to entering the study. A total of 14 hospitalizations occurred during the study but none were related to infection.

Hematology and clinical chemistry parameters were monitored in all subjects prior to each infusion throughout the 12-month period of study. Mean values for all laboratory parameters remained consistent throughout the study period. Three of the hematology values in one subject were outside of the normal range and reported as non-serious adverse experiences that resolved completely. These were a red cell count of  $3.9 \times 10^9/\mu\text{L}$ , hematocrit of 31%, and white cell count of  $3.88 \times 10^9/\mu\text{L}$ . All spontaneously returned to baseline. One subject had an elevated BUN (45 mg/dL) and creatinine (1.4 mg/dL) on one occasion that were reported as non-serious adverse experiences and resolved completely. These values improved to 30 mg/dL and 0.8 mg/dL, respectively, by the next infusion. Six of the patients had a single, transient elevation in serum transaminases. Two additional patients had persistent elevations in transaminases, ALT and AST, which were present at the initiation of the study, prior to the infusion of GAMMAGARD LIQUID. There was no other evidence of liver abnormalities. None of the hematology or chemistry laboratory abnormalities that occurred during the course of the study required clinical intervention and none had clinical consequences.

During the Phase 3 clinical study, viral safety was assessed by serological screening for HBsAg and antibodies to HCV and HIV-1 and HIV-2 prior to, during, and at the end of the study and by Polymerase Chain Reaction (PCR) tests for HBV, HCV, and HIV-1 genomic sequences prior to and at the end of the study. None of the 61 treated subjects were positive prior to study entry and none converted from negative to positive during the 12-month period of study.

#### Postmarketing:

The following is a list of adverse reactions that have been identified and reported during the post-approval use of IGIV products:

<b>Respiratory</b> cyanosis, hypoxemia, pulmonary edema, dyspnea, bronchospasm
<b>Cardiovascular</b> thromboembolism, hypotension
<b>Neurological</b> seizures, tremor
<b>Hematologic</b> hemolysis, positive direct antiglobulin (Coombs) test
<b>General/Body as a Whole</b> pyrexia, rigors
<b>Musculoskeletal</b> back pain
<b>Gastrointestinal</b> hepatic dysfunction, abdominal pain

#### Rare and Uncommon Adverse Events:

<b>Respiratory</b> apnea, Acute Respiratory Distress Syndrome (ARDS), Transfusion Related Acute Lung Injury (TRALI)
<b>Integumentary</b> bullous dermatitis, epidermolysis, erythema multiforme, Stevens-Johnson syndrome
<b>Cardiovascular</b> cardiac arrest, vascular collapse
<b>Neurological</b> coma, loss of consciousness
<b>Hematologic</b> pancytopenia, leukopenia

Because postmarketing reporting of these reactions is voluntary and the at-risk populations are of uncertain size, it is not always possible to reliably estimate the frequency of the reaction to establish a causal relationship to exposure to the product. Such is also the case with literature reports authored independently<sup>46</sup> (see PRECAUTIONS).

#### DOSAGE AND ADMINISTRATION

GAMMAGARD LIQUID should be at room temperature during administration.

Parenteral drug products should be inspected visually for particulate matter and/or discoloration prior to administration. Do not use if particulate matter and/or discoloration is observed. Only clear or slightly opalescent and colorless or pale yellow solutions are to be administered. GAMMAGARD LIQUID should only be administered intravenously. Other routes of administration have not been evaluated. The use of an in-line filter is optional.

For patients with Primary Immunodeficiency, monthly doses of approximately 300 – 600 mg/kg infused at 3 to 4 week intervals are commonly used.<sup>23,24</sup> As there are significant differences in the half-life of IgG among patients with Primary Immunodeficiency, the frequency and amount of immunoglobulin therapy may vary from patient to patient. The proper amount can be determined by monitoring clinical response. The minimum serum concentration of IgG necessary for protection varies among patients and has not been established by controlled clinical studies.

#### Rate of Administration

During the first infusion of the Phase 3 clinical study, GAMMAGARD LIQUID was infused at an initial rate of 0.5 mL/kg/hr (0.8 mg/kg/min). The rate was gradually increased every 30 minutes to a rate of 5.0 mL/kg/hr (8.9 mg/kg/min) if it was well tolerated. However, some patients completed the infusion before the maximum rate could be obtained. During subsequent infusions the initial rate and the rate of escalation were based on their previous infusion history; however, the maximum rate attained during the first infusion was used throughout the remainder of the study. The mean rate attained by all patients was 4.3 mL/kg/hr. Fifty-eight subjects (95%) achieved a maximum rate of 4.0 mL/kg/hr or greater and of these, 16 subjects (26%) attained a rate of 5.0 mL/kg/hr.

In general, it is recommended that patients beginning therapy with IGIV or switching from one IGIV product to another be started at the lower rates and then advanced to the maximal rate if they have tolerated several infusions at intermediate rates of infusion. It is important to individualize rates for each patient.

As noted in the WARNINGS section, **patients who have underlying renal disease or who are judged to be at risk of developing thrombotic events should not be infused rapidly with any IGIV product.** Although there are no prospective studies demonstrating that any concentration or rate of infusion is completely safe, it is believed that risk is decreased at lower rates of infusion.<sup>46</sup> Therefore, as a guideline, it is recommended that these patients who are judged to be at risk of renal dysfunction or thrombotic complications be gradually titrated up to a more conservative maximal rate of less than 3.3 mgIgG/kg/min (< 2 mL/kg/hr).

A rate of administration that is too rapid may cause flushing and changes in pulse rate and blood pressure. Slowing or stopping the infusion usually results in the prompt disappearance of signs. The infusion may then be resumed at a rate that is comfortable for the patient.

#### Drug Interactions

Antibodies in IGIV products may interfere with patient responses to live vaccines, such as those for measles, mumps and rubella.<sup>47,48,49</sup> The immunizing physician should be informed of recent therapy with IGIV products so that appropriate precautions can be taken.

Admixtures of GAMMAGARD LIQUID with other drugs and intravenous solutions have not been evaluated. It is recommended that GAMMAGARD LIQUID be administered separately from other drugs or medications that the patient may be receiving. The product should not be mixed with IGIV products from other manufacturers.

Normal saline should not be used as a diluent. If dilution is preferred, GAMMAGARD LIQUID may be diluted with 5% dextrose in water (D5W).<sup>50</sup> No other drug interactions or compatibilities have been evaluated.

#### HOW SUPPLIED

GAMMAGARD LIQUID is supplied in single use bottles as follows:

NDC Number	Volume	Grams
0944-2700-02	10 mL	1.0
0944-2700-03	25 mL	2.5
0944-2700-04	50 mL	5.0
0944-2700-05	100 mL	10.0
0944-2700-06	200 mL	20.0


## STORAGE

**Refrigeration:** 36 months storage at refrigerated temperature 2° to 8°C (36°-46°F). Do not freeze.

**Room Temperature:** 9 months storage at room temperature 25°C, (77°F) within the first 24 months of the date of manufacture. See below for detailed storage information.

The total storage time of GAMMAGARD LIQUID depends on the point of time the vial is transferred to room temperature. Examples for total storage times are illustrated in Figure 1. The new expiration date must be recorded on the package when the product is transferred to room temperature.

**Figure 1: Storage Guidelines**  
Months from Date of Manufacture



## Storage Details:

- Example 1: If the product is taken out of the refrigerator after 3 months from date of manufacture, it can be stored for 9 months at room temperature. Total storage time is 12 months.
- Example 2: If the product is taken out of the refrigerator after 21 months from the date of manufacture, it can be stored for 3 additional months at room temperature. Total storage time is 24 months.
- After 24 months from date of manufacture, product cannot be stored at room temperature.

To enroll in the confidential, industry-wide Patient Notification System, call 1-888-UPDATE U (1-888-873-2838)

BAXTER and GAMMAGARD LIQUID are trademarks of Baxter International Inc.

**Baxter Healthcare Corporation**  
Westlake Village, CA 91362 USA  
U.S. License No. 140

LE-07-01130

April 2005

## REFERENCES

- Skvaril F. Qualitative and quantitative aspects of IgG subclasses in i.v. immunoglobulin preparations. In: Nydegger UE, ed. Immunotherapy. London: Academic Press; 1981:118-122.
- French M. Serum IgG subclasses in normal adults. Monogr Allergy. 1986;19:100-107.
- Lacy CF, Armstrong LL, Goldman MP, Lance LL. Appendix: Abbreviations and Measurements. Drug Information Handbook. Lexi-Comp; 1999:1254.
- Horowitz B, Prince AM, Hamman J, Watkiewicz C. Viral safety of solvent/detergent-treated blood products. Blood Coagul Fibrinolysis. 1994;5 Suppl 3:S21-S28.
- Kreil TR, Berting A, Kistner O, Kindermann J. West Nile virus and the safety of plasma derivatives: verification of high safety margins, and the validity of predictions based on model virus data. Transfusion. 2003;43:1023-1028.
- Hamamoto Y, Harada S, Kobayashi S, et al. A novel method for removal of human immunodeficiency virus: filtration with porous polymeric membranes. Vox Sang. 1989;56:230-236.
- Yuasa T, Ishikawa G, Manabe S, Sekiguchi S, Takeuchi K, Miyamura T. The particle size of hepatitis C virus estimated by filtration through microporous regenerated cellulose fibre. J Gen Virol. 1991;72 (Pt 8):2021-2024.
- Kempf C, Jentsch P, Poirier B, et al. Virus inactivation during production of intravenous immunoglobulin. Transfusion. 1991;31:423-427.
- Louie RE, Galloway CJ, Dumas ML, Wong MF, Mitra G. Inactivation of hepatitis C virus in low pH intravenous immunoglobulin. Biologics. 1994;22:13-19.
- Golding B. IGIV Clinical Endpoints. Presented at: Blood Products Advisory Committee, 65th Meeting. 17 March 2000. Silver Spring, MD.
- Schiff RL. Intravenous immunoglobulins for treatment of antibody deficiencies. In: Good RA, Lindenlaub E, eds. The Nature, Cellular, and Biochemical Basis and Management of Immunodeficiencies. Symposium Verrier, West Germany. 21-25 September. Stuttgart: F. K. Schattauer Verlag; 1986:523-541.
- Morell A. Pharmacokinetics of intravenous immunoglobulin preparations. In: Lee ML, Strand V, eds. Intravenous Immunoglobulins in Clinical Practice. New York: M. Dekker, Inc.; 1997:1-18.
- Morell A, Skvaril F. Structure and Biological Properties of Immunoglobulins and  $\gamma$ -Globulin Preparations. II. Properties of  $\gamma$ -Globulin Preparations. Schweizerische Medizinische Wochenschrift 1980; 110:80-85.
- Waldmann TA, Strober W. Metabolism of immunoglobulins. Prog Allergy. 1969;13:1-110.
- Stiehm ER. Standard and special human immune serum globulins as therapeutic agents. Pediatrics. 1979;63:301-319.
- Lee ML, Mankariou S, Ochs H, Fischer S, Wedgwood RJ. The pharmacokinetics of total IgG, IgG subclasses, and type specific antibodies in immunodeficient patients. Immunol Invest. 1991;20:193-198.
- Buckley RH. Immunoglobulin replacement therapy: indications and contraindications for use and variable IgG levels achieved. In: Alving BM, Finlayson JS, eds. Immunoglobulins: characteristics and use of intravenous preparations. Washington, D.C.: US Department of Health and Human Services; 1979:3-8.
- Mankariou S, Lee M, Fischer S, et al. The half-lives of IgG subclasses and specific antibodies in patients with primary immunodeficiency who are receiving intravenously administered immunoglobulin. J Lab Clin Med. 1988;112:634-640.
- Pirofsky B. Safety and toxicity of a new serum immunoglobulin G intravenous preparation, IGIV pH 4.25. Rev Infect Dis. 1986;8 Suppl 4:S457-63.
- Pirofsky B. Clinical use of a new pH 4.25 intravenous immunoglobulin preparation (Gamimune-N). J Infect. 1987;15 Suppl 1:29-37.
- Schiff RL. Half-life and clearance of pH 6.8 and pH 4.25 immunoglobulin G intravenous preparations in patients with primary disorders of humoral immunity. Rev Infect Dis. 1986;8 Suppl 4:S449-56.
- Schiff RL, Rudd C. Alterations in the half-life and clearance of IgG during therapy with intravenous gamma-globulin in 16 patients with severe primary humoral immunodeficiency. J Clin Immunol. 1986;6:256-264.
- Eijkhout HW, Der Meer JW, Kallenberg CG, et al. The effect of two different dosages of intravenous immunoglobulin on the incidence of recurrent infections in patients with primary hypogammaglobulinemia. A randomized, double-blind, multicenter crossover trial. Ann Intern Med. 2001;135:165-174.
- Roitman CM, Gelfand EW. Replacement therapy with high dose intravenous gamma-globulin improves chronic sinusopulmonary disease in patients with hypogammaglobulinemia. Pediatr Infect Dis J. 1988;7:S92-S96.
- Balow M, Berger M, Bonilla FA, et al. Pharmacokinetics and tolerability of a new intravenous immunoglobulin preparation, IGIV-C, 10% (Gamunex, 10%). Vox Sang. 2003;84:202-210.
- Ochs HD, Pinciaro PJ, The Octagam Study Group. Octagam(R) 5%, an intravenous IgG product, is efficacious and well tolerated in subjects with primary immunodeficiency diseases. J Clin Immunol. 2004;24:309-314.
- Cayco AV, Perazella MA, Hayslett JP. Renal insufficiency after intravenous immune globulin therapy: a report of two cases and an analysis of the literature. J Am Soc Nephrol. 1997;8:1788-1794.
- Copelan EA, Strohm PL, Kennedy MS, Tutschka PJ. Hemolysis following intravenous immune globulin therapy. Transfusion. 1986;26:410-412.
- Wilson JR, Bhoopalam H, Fisher M. Hemolytic anemia associated with intravenous immunoglobulin. Muscle Nerve. 1997;20:1142-1145.
- Thomas MJ, Misbah SA, Chapel HM, Jones M, Erlington G, Newsom-Davis J. Hemolysis after high-dose intravenous Ig. Blood. 1993;82:3789.
- Kessary-Shoham H, Levy Y, Shoenfeld Y, Lorber M, Gershon H. In vivo administration of intravenous immunoglobulin (IVIg) can lead to enhanced erythrocyte sequestration. J Autoimmun. 1999;13:129-135.
- Rizk A, Gorson KC, Kenney L, Weinstein R. Transfusion-related acute lung injury after the infusion of IVIG. Transfusion. 2001;41:264-268.
- Brannagan TH, III, Nagle KJ, Lange DJ, Rowland LP. Complications of intravenous immune globulin treatment in neurologic disease. Neurology. 1996;47:674-677.
- Dalaks MC. High-dose intravenous immunoglobulin and serum viscosity: risk of precipitating thromboembolic events. Neurology. 1994;44:223-226.
- ElKayam O, Paran D, Milo R, et al. Acute myocardial infarction associated with high dose intravenous immunoglobulin infusion for autoimmune disorders. A study of four cases. Ann Rheum Dis. 2000;59:77-80.
- Gomperts ED, Darr F. Rapid infusion of intravenous immunoglobulin in patients with neuromuscular diseases. Neurology. 2002;58:1444.
- Haplea SS, Farrar JT, Gibson GA, Laskin M, Pizzi LT, Ashbury AK. Thromboembolic events associated with intravenous immunoglobulin therapy [abstract]. Neurology. 1997;48:A54.
- Harkness K, Howell SJ, Davies-Jones GA. Encephalopathy associated with intravenous immunoglobulin treatment for Guillain-Barre syndrome. J Neurol Neurosurg Psychiatry. 1996;60:586.
- Kwan T, Keith P. Stroke following intravenous immunoglobulin infusion in a 28-year-old male with common variable immune deficiency: a case report and literature review. Can J Allergy Clin Immunol. 1999;4:250-253.
- Wolberg AS, Kon RH, Monroe DM, Hoffman M. Coagulation factor XI is a contaminant in intravenous immunoglobulin preparations. Am J Hematol. 2000;65:30-34.
- Woodruff RK, Grigg AF, Firkin FC, Smith IL. Fatal thrombotic events during treatment of autoimmune thrombocytopenia with intravenous immunoglobulin in elderly patients. Lancet. 1986;2:217-218.
- Hammarstrom L, Smith CI. Placental transfer of intravenous immunoglobulin. Lancet. 1986;1:681.
- Morell A, Sidiropoulos D, Herrmann U, et al. IgG subclasses and antibodies to group B streptococci, pneumococci, and tetanus toxoid in preterm neonates after intravenous infusion of immunoglobulin to the mothers. Pediatr Res. 1986;20:933-936.
- Sidiropoulos D, Herrmann U, Jr., Morell A, von Muralit G, Barandun S. Transplacental passage of intravenous immunoglobulin in the last trimester of pregnancy. J Pediatr. 1986;109:505-508.
- Pierce LR, Jain N. Risks associated with the use of intravenous immunoglobulin. Transfusion Med Rev. 2003;17:241-251.
- Tan E, Hajinazarian M, Bay W, Neff J, Mendell JR. Acute renal failure resulting from intravenous immunoglobulin therapy. Arch Neurol. 1993;50:137-139.
- Morbidity and Mortality Weekly Report. Measles, Mumps, and Rubella: Vaccine use and strategies for elimination of measles, rubella, and congenital rubella syndrome and control of mumps. Recommendations for the Advisory Committee on Immunization Practices (ACIP). 98 A.D.;47.
- Peter G. Summary of major changes in the 1994 Red Book: American Academy of Pediatrics. Report of the Committee on Infectious Disease. Pediatrics. 1994;93:1000-1002.
- Siber GR, Werner BG, Halsey NA, et al. Interference of immune globulin with measles and rubella immunization. J Pediatr. 1993;122:204-211.
- Data on file, Baxter Healthcare Corporation.